## Recommended Vaccine Schedule Office of Dr. Manish Dixit. MD

\*Our vaccine schedule differs slightly from the CDC\ACIP recommendation\*

<u>Age</u>	Vaccine(s)	<u>Interval</u>
Birth	Нер В	NA
2 Month	Pediarix, HIB, PCV13, Rota	8 weeks – 6wk minimum
4 Month	Pediarix, HIB, PCV13, Rota	8 weeks – 6wk minimum
6 Month	Pediarix, HIB, PCV13, Rota	8 weeks – 6wk minimum
9 Month	No shots unless catching up or Flu shot – Hgb & Lead screen	
*12 Months	ProQuad	Age dependent – not before birthday
*15 Months	Dtap, HIB, PCV13	6mo interval since last dose of the same
*18 Months	HEP A	May be given as early as 12mo
*24 Months	HEP A	6mo minimum interval between dose 1&2
30 Months	No shots unless catching up or Flu	
4 Years	Kinrix, ProQuad	Age dependent – not before 4 <sup>th</sup> birthday
*10 years	Tdap	Every 10 years here after
11 Years	HPV & Menactra	HPV 3 doses – repeated at 2 & 4 months
16 Years	Menactra & MenB*	MenB is a series - interval dependent on vaccine given

## **Combination Vaccines**

Pediarix: Dtap, Polio, HepB

Kinrix: Dtap, Polio

ProQuad: MMR, Varicella

**Pentacel:** Dtap, Polio, HIB - not presently used in the office

**Specific Dose Intervals** 

HPV(Gardasil) – 3doses - 1st, 2nd in 2mo, 3rd 6mo after 1st dose

MCV4 Menactra – 2 doses, 8wks apart – booster after age 16

MenB **Trumenba** – 3 doses 1st, 2nd in 2mo, 3rd 6mo after 1st dose

MenB **Bexsero** – 2 doses 1 month apart

**Hep A** – at least 6 months apart

**Hep B** – 3 dose minimum - 1st, 2nd in 2mo, 3rd 6mo after 1st dose

**Dtap** – 6week minimum between doses

MenB\* is not yet required - education to be provided and vaccine offered

Call the office 620-421-0002 and ask to speak with the nurse for any specific vaccine questions or inquiries.